A LOCAL CONJECTURE ON BRAUER CHARACTER DEGREES OF FINITE GROUPS

THORSTEN HOLM, WOLFGANG WILLEMS

Abstract. Recently, a new conjecture on the degrees of the irreducible Brauer characters of a finite group was presented in [16]. In this paper we propose a 'local' version of this conjecture for blocks B of finite groups, giving a lower bound for the maximal degree of an irreducible Brauer character belonging to B in terms of the dimension of B and well-known invariants like the defect and the number of irreducible Brauer characters. We also propose a weaker version of this conjecture for which a slight reformulation leads to interesting open questions about traces of Cartan matrices of blocks.

We then show that the strong conjecture is true for blocks with one simple module, blocks of p-solvable groups and blocks with cyclic defect groups. It also holds for many further examples of blocks of sporadic groups, symmetric groups or groups of Lie type. We also show that the weak conjecture is true for blocks of tame representation type.

Introduction

Representation theory of finite groups is an area which is characterized by an enormous number of important long-standing open problems. Many of them deal with ordinary or modular characters, so character theory still is one of the most important and fundamental parts of representation theory. The key aim in this area is to find intimate relations between the numerical invariants given by characters and the structure of the group. Actually, R. Brauer's main motivation for introducing modular characters was to study the structure of finite simple groups.

In this paper we are going to study some of the most important numerical invariants in modular representation theory, namely degrees of irreducible Brauer characters. The main aim is to propose a conjecture relating the degrees of the irreducible characters belonging to a block to some well-known invariants of the block. This can be considered as a block version of a conjecture recently presented in [16] for finite groups. In order to state the conjecture we need to introduce some notation.

Let G be a finite group and let p be a prime dividing the order of G. Moreover, let B be a p-block of G with defect group D where the underlying field of characteristic p is assumed to be algebraically closed. Furthermore let $\mathrm{Irr}_p(B) = \{\varphi_1, \ldots, \varphi_{l(B)}\}$ denote the set of irreducible Brauer characters belonging to B and $\mathrm{Irr}_p(G)$ the set of all irreducible Brauer characters of G.

In [16] the second author proposed the following conjecture relating the degrees of the irreducible Brauer characters of a finite group G to the p'-part of the order of G (i.e. the largest divisor of the order of G not divisible by p).

Global Conjecture [16]: *For any finite group G we have*

$$|G|_{p'} \leq \sum_{\varphi \in \mathrm{Irr}_p(G)} \varphi(1)^2 \tag{1}$$

with equality if and only if the Sylow p-subgroup of G is normal.

Mathematics Subject Classification (2000): Primary: 20C20; Secondary: 15A18, 15A36, 16G60, 20C05
Keywords: Brauer character; Block of finite group; Cartan matrix; Perron-Frobenius eigenvalue.

This conjecture is known to be true for p-solvable groups, groups with a cyclic Sylow p-subgroup, groups of Lie type in the defining characteristic, and asymptotically in many cases in non-defining characteristic, and many more examples (see [16] for details).

The main aim of this paper is to develop a 'local' version of this conjecture, i.e. a version of this conjecture for blocks of finite groups. The crucial aspect is to find a good replacement for the left hand side, which only depends on the block and its invariants, but not on the group to which the block belongs.

As a first suggestion, directly inspired by the above Global Conjecture, we propose the following local version, relating the degrees of the irreducible Brauer characters to well known invariants of the block.

Weak Conjecture: *Let B be a p-block of a finite group, and let D be its defect group. Then we have*

$$\frac{\dim B}{l(B)|D|} \leq \sum_{\varphi \in \mathrm{Irr}_p(B)} \varphi(1)^2 \tag{2}$$

with equality if and only if $l(B) = 1$.

When experimentally computing examples one observes that in most cases the right hand side becomes far larger than the left hand side. So it is tempting to try to improve the conjecture. After having computed and examined an extensive list of examples the best conjecture we propose in this article is the following.

Strong Conjecture: *Let B be a p-block of a finite group, and let D be its defect group. Then we have*

$$\frac{\dim B}{l(B)|D|} \leq \max_{\varphi \in \mathrm{Irr}_p(B)} \{\varphi(1)^2\}. \tag{3}$$

Unfortunately, we could not find a general conjectural characterization for equality in this Strong Conjecture.

If the Strong Conjecture were true then by summing over all blocks of a finite group it had the following immediate 'global' consequence which is a weak version of the Global Conjecture given in [16]. Denote by $l(G)$ the number of p'-conjugacy classes of G (i.e. conjugacy classes of G of elements whose order is not divisible by p). By a classical result of Brauer, $l(G)$ equals the number of irreducible Brauer characters of G.

Weak Global Conjecture: *For any finite group G we have*

$$\frac{|G|_{p'}}{l(G)} \leq \max_{\varphi \in \mathrm{Irr}_p(G)} \{\varphi(1)^2\} \tag{4}$$

with equality if and only if $G = O_p(G) \times O_{p'}(G)$ with $O_{p'}(G)$ abelian.

For $p \neq 2$ we are going to prove in this paper the statement about equality in (4) assuming the Strong Conjecture holds. (Unfortunately, our present proof depends on the classification of finite simple groups.)

At this moment, in either form the conjectures seem to be difficult to prove in general. Still, in this paper we are going to show that the Strong Conjecture is true for various important classes of blocks of finite groups. The results can be summarized as follows.

Theorem 1. *Let B be a p-block of a finite group G with defect group D, and let $l(B)$ be the number of irreducible Brauer characters. Then the Strong Conjecture is true if one of the following holds.*

(a) *$l(B) = 1$, with equality.*

(b) *G is p-solvable.*
Equality holds if and only if the following hold:
(i) *All irreducible Brauer characters of B have the same degree.*
(ii) $\frac{|G|_p}{|D|} = \varphi(1)_p$ *for all* $\varphi \in \mathrm{Irr}_p(B)$.

(c) *D is cyclic.*
Equality holds if and only if the corresponding Brauer tree is a star with exceptional vertex in the center.

Moreover, in addition to these classes of blocks we checked very many further (isolated) examples for which the Strong Conjecture is true, giving further evidence for the validity of the conjecture. Among these examples are many blocks of sporadic simple groups, blocks of (small) symmetric or alternating groups, and certain blocks of groups of Lie type.

For proving the Weak Conjecture we have the following interesting approach via linear algebra. Namely, the following close relation to Cartan matrices will be proved in Section 2.

Proposition 2. *Let B be a p-block of a finite group, and let C be the Cartan matrix of B with trace* $\mathrm{tr}\, C$. *Then the following hold.*

(a) *We have*
$$\frac{\dim B}{\mathrm{tr}\, C} \leq \sum_{\varphi \in \mathrm{Irr}_p(B)} \varphi(1)^2,$$
and equality holds if and only if $l(B) = 1$.

(b) *If $\mathrm{tr}\, C \leq l(B)|D|$, then the Weak Conjecture holds for B.*

This linear algebra approach seem to make the Weak Conjecture more accessible than the Strong Conjecture, since in many cases, the Cartan entries might be known although the degrees of the irreducible characters are not. For instance, we can prove the following result for blocks of tame representation type based on Erdmann's classification of their Cartan matrices.

Theorem 3: *The Weak Conjecture holds for all blocks of tame representation type (i.e. for all 2-blocks with defect group dihedral, semidihedral or generalized quaternion)*

We are not aware of any example of a block such that $\mathrm{tr}\, C \leq l(B)|D|$ does not hold. Actually, we strongly believe that this inequality will be true in general. Note that the inequality is not at all obvious since there are examples of blocks where the diagonal entries in the Cartan matrix are larger than $|D|$. The first example was given by Landrock [10], disproving Problem 22 of Brauer's famous list [1]. Even more striking examples were found by Chastkofsky and Feit [6]; they showed that for the principal 2-blocks of the Suzuki groups $\mathrm{Suz}(2^m)$ the first diagonal Cartan entry is asymptotically $|D|^{3/2}$.

Further evidence for the validity of our global conjectures is provided by analogies between degrees of irreducible characters and lengths of conjugacy classes. Such analogies have been observed by several authors (see for instance [8]). The statements about conjugacy class lengths which correspond to our Global Conjecture and Weak Global Conjecture on character degrees can actually be proved. We plan to address this issue in a separate note.

1. Discussions around the conjectures

In this section we collect some of the evidence we have supporting the conjectures, and discuss various aspects around the conjectures. In particular, we indicate that the Strong Conjecture is true for blocks with one simple module and for blocks of p-solvable groups.

1.1 Blocks with one simple module. Let B be a block with one simple module, i.e. $l(B) = 1$. Clearly, in this case the Weak and the Strong Conjecture coincide. The validity of the conjectures

in this case is a classical result of R. Brauer, stating that in the case $l(B) = 1$ equality holds in (2) (see [2], Theorems 2 and 4). This proves Theorem 1(i).

For an arbitrary block B, Brauer showed in [2] that the p-part of $\frac{\dim B}{|D|}$ is equal to the p-part of $\sum_{\varphi \in \mathrm{Irr}_p(B)} \varphi(1)^2$. Thus equality in (2) forces $p \nmid l(B)$. But the 'only if' part of the characterization of equality remains open.

1.2 It might be tempting to try to improve the left hand side of the inequality in the Weak Conjecture. We shall see below that for some interesting classes of blocks one can actually show that

$$\frac{\dim B}{|D|} \leq \sum_{\varphi \in \mathrm{Irr}_p(B)} \varphi(1)^2, \tag{5}$$

which then implies that the Strong Conjecture holds for B.

But the following examples show that the factor $l(B)$ in the Weak Conjecture can in general not be dropped and not be replaced by any constant.

Let $G = \mathrm{A}_5$ denote the alternating group on 5 letters and let B be the principal 2-block of G. Then B contains three irreducible Brauer characters of degrees $1, 2$ and 2, respectively. The corresponding projective characters are of degree $12, 8, 8$. Thus we obtain

$$\frac{\dim B}{|D|} = \frac{44}{4} = 11 > 9 = 1 + 4 + 4 = \sum_{\varphi \in \mathrm{Irr}_2(B)} \varphi(1)^2.$$

Now we take n copies of $G = \mathrm{A}_5$ and consider the principal 2-block $B^{(n)}$ of the direct product $G^{(n)} = G \times \ldots \times G$. Clearly $B^{(n)}$ has defect group $D^{(n)} = D \times \ldots \times D$. It is not hard to compute that

$$\frac{\dim B^{(n)}}{|D^{(n)}|} = 11^n > 9^n = \sum_{\varphi \in \mathrm{Irr}_2(B^{(n)})} \varphi(1)^2.$$

Thus the factor $l(B)$ in the weak conjecture can not be replaced by a constant.

Similar examples are provided by the maximal non-principal 3-block of $6\mathrm{A}_7$, and by the principal 2-block of $\mathrm{SL}(2, 5)$, showing that the above phenomenon neither depends on the prime 2 nor on D being abelian.

Note that the above 2-blocks $B^{(n)}$ of $G^{(n)} = A_5 \times \ldots \times A_5$ nicely satisfy our Strong Conjecture. In fact, we have

$$\frac{\dim B^{(n)}}{l(B^{(n)})|D^{(n)}|} = \left(\frac{11}{3}\right)^n < 2^{2n} = \max_{\varphi \in \mathrm{Irr}_2(B^{(n)})} \{\varphi(1)^2\}.$$

1.3 We would like to mention that equality in (5) does not imply $l(B) = 1$ as the following example shows.

Let G be a finite group with a normal Sylow p-subgroup P. Furthermore let $C_G(P) \leq P$. Thus G is a p-solvable group with exactly one p-block (see [5], Chap. V, 3.11). Since P is normal in G it is contained in the kernel of any irreducible representation in characteristic p . Thus we have $p \nmid \varphi(1)$ for all $\varphi \in \mathrm{Irr}_p(G)$. A result of Fong (see [5], Chap. X, 3.2) yields $\Phi(1) = |P|\varphi(1)$ where Φ denotes the projective character corresponding to φ. Thus equality holds in (5), but $l(B) > 1$ if $P \neq G$.

1.4 Blocks of p-solvable groups. Let B be a block of a p-solvable group G. We are going to show that the inequality (5) holds for B, which implies the Strong Conjecture for blocks of p-solvable groups.

Let Φ_φ denote the projective character corresponding to the irreducible Brauer character φ. Again by Fong's result we have $\Phi_\varphi(1) = |G|_p\varphi(1)_{p'}$ (see [5], Chap. X, 3.2) and, by a result of Green, we have $\frac{|G|_p}{|D|} \mid \varphi(1)$ (see [5], Chap. IV, 4.5). Thus we get

$$\frac{\dim B}{|D|} = \frac{1}{|D|}\sum_{\varphi\in\mathrm{Irr}_p(B)}\varphi(1)\Phi_\varphi(1) = \frac{|G|_p}{|D|}\sum_{\varphi\in\mathrm{Irr}_p(B)}\varphi(1)\varphi(1)_{p'} \le \sum_{\varphi\in\mathrm{Irr}_p(B)}\varphi(1)^2.$$

For proving Theorem 1(ii) it remains to characterize equality in our Strong Conjecture. From the discussions above we have

$$\frac{\dim B}{|D|} \le \sum_{\varphi\in\mathrm{Irr}_p(B)}\varphi(1)^2 \le l(B)\max_{\varphi\in\mathrm{Irr}_p(B)}\{\varphi(1)^2\}$$

where the first inequality is an equality if and only if $\frac{|G|_p}{|D|} = \varphi(1)_p$ for all $\varphi \in \mathrm{Irr}_p(B)$. Clearly, the second inequality becomes an equality if and only if all irreducible Brauer characters have the same degree. This proves Theorem 1(ii).

1.5 Connections to classical open problems. Let $k(B)$ denote the number of ordinary irreducible characters in an arbitrary block B, with defect group D. Clearly $l(B) \le k(B)$. Problem 20 of Brauer's famous list [1] asks whether always $k(B) \le |D|$. If we assume that this has an affirmative answer we get $l(B)|D| \le |D|^2$ and our Weak Conjecture leads to

$$\frac{\dim B}{|D|^2} \le \sum_{\varphi\in\mathrm{Irr}_p(B)}\varphi(1)^2. \tag{6}$$

This obviously holds true if

$$\Phi_\varphi(1) \le |D|^2\varphi(1) \quad \text{for all } \varphi \in \mathrm{Irr}_p(B) \tag{7}$$

where Φ_φ is the projective character corresponding to φ. Note that in general $\Phi_\varphi(1)$ is larger than $l(B)|D|$ (for instance, take the projective character corresponding to the trivial module for the smallest Suzuki group in characteristic 2). We are not aware of any example for which (7) does not hold.

1.6 What are the connections between the Global Conjecture from [16] and our Weak Conjecture? An affirmative answer to the Weak Conjecture would also prove a weaker version of the Global Conjecture. In fact, if $l(G)$ denotes the number of p'-conjugacy classes of G then by summing up (2) over all blocks B of G we would get

$$\frac{|G|_{p'}}{l(G)} \le \sum_{\varphi\in\mathrm{Irr}_p(G)}\varphi(1)^2 \tag{8}$$

with equality if and only if G is a p-group. In fact, suppose equality holds. This means that G has only one block with exactly one irreducible Brauer character. Thus the trivial character is the only irreducible Brauer character and therefore G has only one conjugacy class of p'-elements. This shows that G is a p-group. On the other hand if G is a p-group then obviously equality holds in (8).

(Note that in general $\Phi_\varphi(1) \not\le |P|l(G)\varphi(1)$ where P is a Sylow p-subgroup of G; take the projective character corresponding to the trivial character for the smallest Suzuki group in characteristic 2.)

1.7 There is a similar direct connection between our Strong Conjecture and the Global Conjecture. If the Strong Conjecture holds then by summing over all blocks of a finite group G we would get that

$$\frac{|G|_{p'}}{l(G)} \le \max_{\varphi\in\mathrm{Irr}_p(G)}\{\varphi(1)^2\} \tag{9}$$

which is also a weak version of the Global Conjecture (though clearly stronger than (8)). Moreover, if $p \neq 2$ then equality holds in (9) if and only if $G = O_p(G) \times O_{p'}(G)$ with abelian $O_{p'}(G)$.

In fact, if G has this structure then equality holds in (9) since $|G|_{p'} = |O_{p'}(G)| = l(G)$ (all equal to the number of p'-conjugacy classes of G) and all irreducible Brauer characters are of degree 1. In order to prove the converse suppose that the Strong Conjecture holds and that equality holds in (9). First we prove by induction that G is p-solvable. We put $H = O_{p'}(G)$ and $\overline{G} = G/H$. Since the group algebra of $\overline{G}$ is a direct summand of the group algebra of G over the underlying field one can show that

$$\frac{|\overline{G}|_{p'}}{l(\overline{G})} = \max_{\varphi \in \mathrm{Irr}_p(\overline{G})} \{\varphi(1)^2\}.$$

If $H \neq \langle 1 \rangle$, then by induction $\overline{G}$ is p-solvable. Hence, G is p-solvable. Next we assume that $O_p(G) \neq \langle 1 \rangle$. If $\hat{G} = G/O_p(G)$ then equality in (9) implies that

$$\frac{|\hat{G}|_{p'}}{l(\hat{G})} = \max_{\varphi \in \mathrm{Irr}_p(\hat{G})} \{\varphi(1)^2\}$$

and the induction argument yields again that $\hat{G}$ is p-solvable. Therefore, also G is p-solvable. Finally, we may assume that G has a minimal normal subgroup, say N, which is a direct product of isomorphic non-abelian simple groups whose orders are divisible by p. The equality in (9) implies that all blocks of G are of maximal defect. Thus, by Knörr's theorem (see [12], 9.26), all blocks of N are of maximal defect. By a result of Michler (see [11], Theorem 5.3), a finite non-abelian simple group has blocks of different defects for $p \neq 2$. Thus we have a contradiction in this case and the claim that G is p-solvable has been proved.

Next we show that $G = O_p(G) \times O_{p'}(G)$ with abelian $O_{p'}(G)$. Since

$$|G|_{p'} = \sum_{\varphi \in \mathrm{Irr}_p(G)} \varphi(1)_{p'}\varphi(1) \leq l(G) \max_{\varphi \in \mathrm{Irr}_p(G)} \{\varphi(1)^2\} = |G|_{p'}$$

all φ are of degree 1. In particular

$$G' \leq \cap_{\varphi \in \mathrm{Irr}_p(G)} \mathrm{Ker}\, \varphi = O_p(G)$$

where G' denotes the commutator subgroup of G. Thus G has a normal Sylow p-subgroup with abelian factor group. The equation

$$\frac{|G|_{p'}}{l(G)} = \max_{\varphi \in \mathrm{Irr}_p(G)} \{\varphi(1)^2\} = 1$$

shows that all p'-conjugacy classes have length 1. Thus a Hall p'-subgroup of G is central in G which proves the assertion.

2. Eigenvalues and Brauer characters

Associated to any p-block B is the Cartan matrix $C = C_B$. If $< \cdot, \cdot >$ denotes the Euclidean inner product on $\mathbb{R}^l$ and $\overline{\varphi} = (\varphi_1(1), \ldots, \varphi_l(1))$ the dimension vector then the dimension of the block (as a vector space over the ground field) can be expressed as $\dim B = < C\overline{\varphi}, \overline{\varphi} >$. If $\rho(C)$ denotes the Perron-Frobenius eigenvalue of C, i.e.

$$\rho(C) = \max\{|\lambda| \mid \lambda \text{ is an eigenvalue of } C\},$$

then, since C is a Hermitian matrix,

$$\frac{< C\overline{\varphi}, \overline{\varphi} >}{< \overline{\varphi}, \overline{\varphi} >} \leq \rho(C).$$

In particular we have the following result.

Proposition 2.1. (Kiyota-Wada [9])

$$\frac{\dim B}{\rho(C)} = \frac{< C\overline{\varphi}, \overline{\varphi} >}{\rho(C)} \leq < \overline{\varphi}, \overline{\varphi} > = \sum_{\varphi \in \mathrm{Irr}_p(B)} \varphi(1)^2. \tag{10}$$

In order to prove our Weak Conjecture it therefore suffices to show that $\rho(C) \leq l(B)|D|$ holds for any block. Usually, $\rho(C)$ is hard to determine and there is no general upper bound known which only depends on invariants of B, and not on the group. Moreover equality in (10) does not seem to be easy to characterize. In fact, suppose equality holds in (10). Since C is Hermitian this implies $C\overline{\varphi} = \rho(C)\overline{\varphi}$. Thus if Φ_φ denotes the projective character corresponding to φ then we have $\Phi_\varphi(1) = \rho(C)\varphi(1)$. By the results of Brauer [2] we obtain that $\rho(C) = |D|$. On the other hand if $\Phi_\varphi(1) = |D|\varphi(1)$ for all $\varphi \in \mathrm{Irr}_p(B)$ then equality holds in (10).

Such situations occur for all blocks if G is p-solvable of p-length 1. In particular, equality in (10) does not imply $l(B) = 1$ in general.

Despite the difficulties in general, for blocks of p-solvable groups we have the following characterization of equality in (10).

Proposition 2.2. *Let G be a p-solvable group. For the principal p-block B the following statements are equivalent.*

(a) $\frac{\dim B}{\rho(C)} = \sum_{\varphi \in \mathrm{Irr}_p(B)} \varphi(1)^2$, *i.e. equality holds in (10).*

(b) *G is of p-length* 1.

Proof: As mentioned above, (b) implies (a) (see [13], Satz 4.2). To prove the other direction let P denote a Sylow p-subgroup of G. By the remarks preceding the proposition we have $\Phi_\varphi(1) = |P|\varphi(1)$ for all $\varphi \in \mathrm{Irr}_p(B)$. This implies $\Phi_\varphi = \varphi\Phi_1$ where Φ_1 corresponds to the trivial character. If $p \mid \varphi(1)$ then the multiplicity of the trivial character in $\varphi\overline{\varphi}$ is at least 2 (see [7], Chap. VII, 8.5 d)). Thus

$$2 \leq (\Phi_1, \varphi\overline{\varphi}) = (\Phi_1\varphi, \varphi) = (\Phi_\varphi, \varphi) = 1,$$

a contradiction. So all irreducible Brauer characters in B are of p'-degree. Since G is assumed to be p-solvable and B is the principal block we conclude that B is isomorphic to the group algebra of $G/O_{p'}(G)$. The condition $p \nmid \varphi(1)$ for all φ in B implies that $G/O_{p'}(G)$ has a normal Sylow p-subgroup (a well-known result of Wallace, see [15] or [3]). Thus G is of p-length 1.

Our next aim is to establish a linear algebra criterion for proving the Weak Conjecture, namely Proposition 2 as stated in the introduction. We begin by recalling some useful fundamental facts about Cartan matrices of blocks.

Lemma 2.3. *The Cartan matrix of any p-block B is positive definite.*

Proof: If D denotes the decomposition matrix of B then $C = D^T D$. Then we have $(Cx, x) = (D^T Dx, x) = (Dx, Dx)$ for all $x \in \mathbb{R}^l$ where $l = l(B)$. If $(Dx, Dx) = 0$ then $Dx = 0$ and therefore $Cx = 0$. But C is regular, hence $x = 0$. This proves that $(Cx, x) > 0$ for all $0 \neq x \in \mathbb{R}^l$.

Corollary 2.4. *If C is the Cartan matrix of a p-block then*

$$\rho(C) \leq \mathrm{tr}\, C.$$

Proof: Since C is positive definite by Lemma 2.3, all eigenvalues of C are positive and they sum up to the trace of C. In particular $\rho(C) \leq \mathrm{tr}\, C$.

Proposition 2.5. *Let B be a p-block of a finite group, with Cartan matrix C. Then*

$$\frac{\dim B}{\mathrm{tr}\, C} \leq \sum_{\varphi \in \mathrm{Irr}_p(B)} \varphi(1)^2,$$

and equality holds if and only if $l(B) = 1$.

Proof: By Proposition 2.1 and Corollary 2.4 we have

$$\frac{\dim B}{\operatorname{tr} C} \leq \frac{\dim B}{\rho(C)} \leq \sum_{\varphi \in \operatorname{Irr}_p(B)} \varphi(1)^2.$$

Suppose equality holds, i.e. $\operatorname{tr} C = \rho(C)$. Since all eigenvalues are positive, and their sum is $\operatorname{tr} C$, we deduce that $l(B) = 1$.

The above proposition gives a possible linear algebra approach to proving the Weak Conjecture provided the following question has an affirmative answer.

Question 2.6. *Let B be a p-block of a finite group, with defect group D and Cartan matrix C. Is it true that*

$$\operatorname{tr} C \leq l(B)|D|? \tag{11}$$

Since C is positive definite we have

$$\det C \leq \prod_{\varphi \in \operatorname{Irr}_p(B)} c_{\varphi\varphi},$$

hence

$$l\sqrt[l]{\det C} \leq l\sqrt[l]{\prod_{\varphi \in \operatorname{Irr}_p(B)} c_{\varphi\varphi}} \leq \sum_{\varphi \in \operatorname{Irr}_p(B)} c_{\varphi\varphi} = \operatorname{tr} C.$$

Using the elementary divisors of C we see that $\sqrt[l]{\det C} \leq |D|$ in accordance with (11).

3. Blocks with cyclic defect groups

This section deals with blocks with cyclic defect group and contains a proof of our Strong Conjecture for these blocks (i.e. a proof of Theorem 1 (iii)). The proof is very much based on the well known structure theory of these blocks, and it also follows from work of Kiyota and Wada [9]. For the sake of completeness and for the convenience of the reader we include a direct proof here. We actually show the following stronger statement.

Proposition 3.1. *For a p-block B with cyclic defect group D we have*

$$\frac{\dim B}{|D|} \leq \sum_{\varphi \in \operatorname{Irr}(B)} \varphi(1)^2 \tag{12}$$

with equality if and only if the corresponding Brauer tree of B is a star with exceptional vertex, if there is one, in the center.

Proof: Let e denote the inertial index of B. In other words e is the number $l(B)$ of irreducible Brauer characters in B. If we put $m = \frac{|D|-1}{e}$ then the shape of the Brauer tree yields

$$0 \leq C = (c_{i,j}) \leq \begin{pmatrix} m+1 & & \\ & \ddots & \\ m & & m \\ & & m+1 \end{pmatrix}$$

for the Cartan matrix C of B where $\leq$ for matrices means $\leq$ in each position. If we put $\overline{\varphi} = (\varphi_1(1), \ldots, \varphi_e(1))$ and $\overline{1} = (1, \ldots, 1)$ then by the Cauchy-Schwarz inequality

$$| < \overline{\varphi}, \overline{1} > |^2 \leq < \overline{\varphi}, \overline{\varphi} > < \overline{1}, \overline{1} >,$$

hence

$$(*) \qquad (\sum_{i=1}^{e} \varphi_i(1))^2 \leq e \sum_{i=1}^{e} \varphi(1)^2.$$

Thus we get

$$\begin{aligned}
\dim B &= \textstyle\sum_{i=1}^{e} \varphi_i(1)\Phi_i(1) \\
&= \textstyle\sum_{i,j=1}^{e} \varphi_i(1)c_{i,j}\varphi_j(1) \\
&\leq m\textstyle\sum_{i,j=1}^{e} \varphi_i(1)\varphi_j(1) + \sum_{i=1}^{e} \varphi_i(1)^2 \\
&= m\textstyle\sum_{i=1}^{e} \varphi_i(1)\sum_{j=1}^{e} \varphi_j(1) + \sum_{i=1}^{e} \varphi_i(1)^2 \\
&\leq me\textstyle\sum_{i=1}^{e} \varphi_i(1)^2 + \sum_{i=1}^{e} \varphi_i(1)^2 \qquad \text{by } (*) \\
&= (me+1)\textstyle\sum_{i=1}^{e} \varphi_i(1)^2 \\
&= |D|\textstyle\sum_{i=1}^{e} \varphi_i(1)^2.
\end{aligned}$$

Suppose equality holds in (12). It follows that

$$C = (c_{i,j}) = \begin{pmatrix} m+1 & & \\ m & \ddots & m \\ & & m+1 \end{pmatrix}.$$

This means that the Brauer tree is a star with the exceptional vertex (if there is one) in the center. Thus all the degrees $\varphi_i(1)$ are equal.

4. Cartan matrices of tame blocks

In this section we give a proof of our Weak Conjecture for blocks of tame representation type, i.e. for all 2-blocks of finite groups having a dihedral, semidihedral or generalized quaternion defect group. For the proof we use the linear algebra approach outlined in Section 2, namely we are going to show that the trace of the Cartan matrix can be bounded above in terms of the order of the defect group. The Cartan matrices of tame blocks are known due to the extraordinary classification of K. Erdmann of tame blocks up to Morita equivalence [4].

It seems to be hard to prove our Strong Conjecture for tame blocks since the degrees of irreducible Brauer characters for an arbitrary tame block are not known in general.

Blocks of tame representation type have at most three simple modules. Since our conjectures are true for blocks with one simple module we can restrict our attention to the blocks with two or three simple modules. We refer to [4] for background on tame blocks, and also for details and notations.

Proposition 4.1. *Let B be a block of a finite group of tame representation type, with defect group D. Then for the trace of the Cartan matrix C of B the following holds.*

(a) *If B has two simple modules, then we have* $\operatorname{tr} C \leq \frac{5}{4}|D| + 2$.

(b) *If B has three simple modules then we have* $\operatorname{tr} C \leq \frac{3}{2}|D| + 4$.

Proof: (a) From Erdmann's classification we see that such a block B has a Cartan matrix of the form

$$\begin{pmatrix} 4k & 2k \\ 2k & k+r \end{pmatrix}$$

with natural numbers k and r where $\{k, r\} = \{1, \frac{|D|}{4}\}$ or $\{k, r\} = \{2, \frac{|D|}{4}\}$. For any possiblity of k and r we obtain that $\operatorname{tr} C \leq \frac{5}{4}|D| + 2$ (note that $|D| \geq 8$ since a block with Klein four defect group can not have two simple modules).

(b) There are various families of algebras in Erdmann's classification which (may) contain blocks of finite groups. They are denoted $D(3\mathcal{A})_1$, $D(3\mathcal{B})_1$ and $D(3\mathcal{K})$ in the dihedral case, $SD(3\mathcal{A})_1$, $SD(3\mathcal{B})_1$, $SD(3\mathcal{C})_2$ and $SD(3\mathcal{D})$ in the semidihedral case, and $Q(3\mathcal{A})_2$, $Q(3\mathcal{B})$ and $Q(3\mathcal{K})$ in the quaternion case.

The families $D(3\mathcal{A})_1$ and $Q(3\mathcal{A})_2$ have Cartan matrices of the form

$$\begin{pmatrix} 4k & 2k & 2k \\ 2k & k+a & k \\ 2k & k & k+a \end{pmatrix}$$

where $k \in \mathbb{N}$ and $a \in \{1,2\}$. Blocks of finite groups only occur for $k = \frac{|D|}{4}$. In particular, $\operatorname{tr} C \leq \frac{3}{2}|D| + 4$.

The algebras $D(3\mathcal{K})$ and $Q(3\mathcal{K})$ have Cartan matrices of the form

$$\begin{pmatrix} 2a & a & a \\ a & k+a & k \\ a & k & k+a \end{pmatrix}$$

where $k \in \mathbb{N}$ and $a \in \{1,2\}$. For blocks of finite groups we have $k = \frac{|D|}{4}$. In particular $\operatorname{tr} C \leq \frac{|D|}{2} + 8 \leq \frac{3}{2}|D| + 4$ (since $|D| \geq 4$ for tame blocks).

The algebras in the families $SD(3\mathcal{B})_1$, $SD(3\mathcal{D})$ and $Q(3\mathcal{B})$ all have Cartan matrices of the form

$$\begin{pmatrix} 4 & 2 & 2 \\ 2 & s+1 & 1 \\ 2 & 1 & 3 \end{pmatrix}$$

for a natural number $s \geq 2$. Blocks of finite groups only occur for $s = \frac{|D|}{4} \geq 2$. In particular we get that $\operatorname{tr} C \leq \frac{|D|}{4} + 8 \leq \frac{3}{2}|D| + 4$.

The algebras in the family $D(3\mathcal{B})_1$ have Cartan matrix of the form

$$\begin{pmatrix} 4 & 2 & 2 \\ 2 & s+1 & 1 \\ 2 & 1 & 2 \end{pmatrix}$$

for some $s \geq 1$. Blocks of finite groups occur for the parameter $s = \frac{|D|}{4}$, hence $\operatorname{tr} C \leq \frac{|D|}{4} + 7$ for blocks.

The algebras in the family $SD(3\mathcal{A})_1$ have Cartan matrix of the form

$$\begin{pmatrix} 4k & 2k & 2k \\ 2k & k+1 & k \\ 2k & k & k+2 \end{pmatrix}$$

for some natural number k. Blocks of finite groups occur for parameters $k = \frac{|D|}{4}$, i.e. $\operatorname{tr} C \leq \frac{3}{2}|D| + 3$ for blocks in this family.

There are two more families for which it is still an open question whether blocks of finite groups actually occur in these families.

The Cartan matrices of algebras in the family $SD(3\mathcal{H})$ are of the form

$$\begin{pmatrix} 3 & 2 & 1 \\ 2 & s+2 & s \\ 1 & s & s+1 \end{pmatrix}$$

for some $s \geq 2$. If blocks occur, then they occur for $s = \frac{|D|}{4}$. In particular, for blocks we have $\operatorname{tr} C \leq \frac{|D|}{2} + 5 \leq \frac{3}{2}|D| + 4$.

The Cartan matrices for algebras in the family $SD(3\mathcal{C})$ are of the form

$$\begin{pmatrix} k+s & k & k \\ k & k+1 & k-1 \\ k & k-1 & k+1 \end{pmatrix}$$

If blocks of finite groups occur then we have $\{k, s\} = \{2, \frac{|D|}{4}\}$. For $k = 2$, $s = \frac{|D|}{4}$ we get $\operatorname{tr} C \leq \frac{|D|}{4} + 8$, in the other case we get $\operatorname{tr} C \leq \frac{3}{4}|D| + 4$.

This completes the proof of the proposition.

As an immediate consequence of the above proposition, we obtain that if B is a block of finite group with tame representation type then $\operatorname{tr} C \leq l(B)|D|$ where D is the defect group. (Note that a block with a Klein four group as defect group has three irreducible Brauer characters.) By Proposition 2.5 we can conclude that the Weak Conjecture holds for all tame blocks, thus proving Theorem 3.

References

[1] R. Brauer, *Representations of finite groups.* Lectures on modern Mathematics, John Wiley, New York 1963.

[2] R. Brauer, *Notes on representations of finite groups, I.* J. London Math. Soc. (2), 13 (1976), 162-166.

[3] P. Brockhaus *On the radical of a group algebra.* J. Algebra **95** (1985), 454-472.

[4] K. Erdmann, *Blocks of Tame Representation Type and Related Algebras.* Lecture Notes in Mathematics **1428**, Springer (1990).

[5] W. Feit, *The representation theory of finite groups.* North Holland, Amsterdam 1982.

[6] L. Chastkofsky, W. Feit, *On the projective characters in characteristic 2 of the groups Suz*(2^m) *and* $Sp_4(2^n)$. Inst. Hautes Études Sci. Publ. Math. No. 51 (1980), 9–35.

[7] B. Huppert, N. Blackburn, *Finite groups II, III* Springer Verlag, Berlin, Heidelberg, New York, 1982.

[8] B. Huppert, *Research in representation theory at Mainz (1984-1990).* In: G.O. Michler, C.M. Ringel (eds.), Representation theory of finite groups and finite-dimensional algebras. Birkhäuser, Basel, 1991.

[9] M. Kiyota, T. Wada, *Some remarks on eigenvalues of the Cartan matrix in finite groups.* Comm. Algebra **21(11)** (1993), 3839-3860.

[10] P. Landrock, *A counterexample to a conjecture on the Cartan invariants of a group algebra.* Bull. London Math. Soc. **5** (1973), 223-224.

[11] G. Michler, *A finite simple group of Lie type has p-blocks of different defects.* J. Algebra **104** (1986), 220-230.

[12] G. Navarro, *Characters and Blocks of Finite Groups.* Lecture Notes Series **250**, Cambridge University Press, Cambridge 1998.

[13] W. Schwarz, *Die Struktur modularer Gruppenringe endlicher Gruppen der p-Länge* 1. J. Algebra **60** (1979), 51-75.

[14] T. Wada, *A lower bound of the Perron-Frobenius eigenvalue of the Cartan matrix for a finite group.* Arch. Math. **73** (1999), 407-413.

[15] D.A.R. Wallace, *On the radical of a group algebra.* Proc. Amer. Math. Soc. **12** (1961), 133-137.

[16] W. Willems, *On Degrees of Irreducible Brauer Characters.* Trans. Am. Math. Soc., to appear.

Thorsten Holm
Department of Pure Mathematics, University of Leeds
Leeds LS2 9JT, United Kingdom
E-mail address: tholm@maths.leeds.ac.uk

Wolfgang Willems
Otto-von-Guericke-Universität, Institut für Algebra und Geometrie
Postfach 4120, 39016 Magdeburg, Germany
E-mail address: wolfgang.willems@mathematik.uni-magdeburg.de